Tabel met studiekarakteristieken - Continuïteit van zorg

Eerste auteur, jaartal	**Inhoud**					
	Studiedesign	Setting, land	Populatie	Geestelijk verzorger	Verpleegkundige	Resultaten
Wilkie, 2024[2]	Stepped-wedge cluster RCT	Amerika, polikliniek	579 volwassen kankerpatiënten in poliklinische palliatieve zorg, ≥55 jaar (gemiddeld 66,4 jaar), 59% vrouw	Waardigheids- therapie volgens handleiding. Vier contactmomenten uitgevoerd door geestelijk verzorger	Waardigheids-therapie volgens handleiding. Vier contactmomenten uitgevoerd verpleegkundige	Geen significant verschil tussen waardigheidstherapie geleid door geestelijk verzorger en verpleegkundige gevonden op de DIS (p = 0.48).

Referenties

1. Scarton, L. et al. Dignity Impact as a Primary Outcome Measure for Dignity Therapy. Am J Hosp Palliat Care 35, 1417–1420 (2018).
2. Wilkie, D. J. et al. Engaging Mortality: Effective Implementation of Dignity Therapy. J Palliat Med 27, 176–184 (2024).
3. Martinuz, M. et al. Do you want some spiritual support? Different rates of positive response to chaplains' versus nurses' offer. J Pastoral Care Counsel 67, 3 (2013).