# **Defining patients eligible for palliative care** by GPs and community nurses

Influence of implementation of PaTz palliative home care groups

Bregje Onwuteaka-Philipsen, Annicka van der Plas, Bart Schweitzer, Roeline Pasman Department of public and occupational health, VU University medical center, Amsterdam





# What is a PaTz group?

In PaTz groups (inspired on GSF) general practitioners (GPs) and community nurses in a neighbourhood:

- meet 6x per year in presence of a palliative care consultant
- identify patients in need of palliative care
- discuss palliative care needs and options

# Identifying patients with palliative care needs

For identifying patients in need of palliative care

To study whether the implementation of PaTz had an effect on elements PaTz participants mention in defining who is eligible for palliative care

### Results

- mentioned in defining patients eligible for

# Methods


- Questionnaire study before and 14 months after start of PaTz group
- Population: participants of 40 PaTz groups:

GPs: pre test 235 (65%), post test 197 (52%)

Community nurses: pre test 54 (54%), post test 58 (52%)

- Open ended question on giving a definition of patients eligible for palliative care
- Coding of elements of definition in categories
- Selection of respondents who answered the definition question pre and post (n=164)

# Five most mentioned elements of definition by all PaTz participants in pre test and post test



\* Significant difference between pre test and post test (mcNemar test, p<0.05)

### Conclusion

- PaTz resulted in a positive change with regard to defining who is eligible for palliative care
- This is potentially due to the attention for the surprise question in the PaTz method
- This is important to achieve timely identification of patients with palliative care needs



